



A Stochastic Model for Rice Yields Forecast

M. Yahaya* and A. Lawal

Department of Mathematics, Federal University of Technology Minna, Nigeria

ABSTRACT

Rice (*Oryza sativa*) is one of most consumed food crops in Nigeria and in around the world. Demand for rice has been increasing at fast rate in Nigeria due to population growth and the banning of its importation in to the country has further led to its shortage in supply. This research is poised to provide necessary information to the government to boost rice production in the Nigeria. In view of this, a stochastic mathematical model that makes forecast of rice yields with respect to climatic elements (Rainfall and Temperature) has been developed and implemented in Niger State. The relevant data used in this research were collected from the Niger state Bureau of statistics over a period of 16years (2001-2016). Validity test for the model was conducted which indicates 80% accuracy, this shows that the model is reliable and dependable. Both the validity test and the future forecast of the model showed prevalence of high rice yields, this shows that Niger State is a better place for rice farming. Therefore, results from study is an important information to Federal Government to boost local rice production in country.

Keywords: Hidden Markov Model, Stochastic Model, Rice, Niger State, Baum-Welch Algorithm

INTRODUCTION

The banning of importation of rice into the country coupled with population growth has led to severe shortage in its supply. Hence, this research is aimed at developing a stochastic model for rice yields forecast with a view of providing necessary information to the government to boost local rice production in the country. Rice is the staple food for more than three billion people worldwide. Rice area accounted for approximately 11.5% of the world's arable land area and it provides almost 19% of the global dietary energy in recent times and its annual average consumption per capita is about 65 kg (Ajithet *al.*, 2017). Therefore, rice area mapping and forecasting is vital for food security, where demands often exceed production due to an ever-increasing population. The aim of yield forecasting, is to give a precise, scientifically sound and independent forecasts of crops' yield as early as possible during the crops' growing season by considering the effect of the weather and climate (Basso, 2013). Timely and accurate estimation of rice areas and forecasting can provide important information for governments, planners, and decision makers in formulating policies with regard to import or export and in the event of shortfall or surplus (Mostafa, 2015).

Rice (*Oryza sativa* L.) is a primary food for more than three billion people worldwide (Khush, 2005). It is cultivated on about 11.5% of the world's arable land during 2012, however more than 88% production is observed in Asian countries. In the recent decades, two major issues like population growth in particular to the rice consuming countries where ~60% of the world's population lives and climate change put enormous pressure on the rice demand (IPCC, 2014). As such, understanding (in other words, forecasting) the amount of rice production prior to the end of growing season is critical in order to ensure food security (Bastiaanssen & Ali, 2003; Nuarsaet *al.*, 2012; Huang *et al.*, 2013; Son *et al.*, 2013). This sort of forecasting may help the governments, planners, and decision makers to formulate appropriate policies to: quantify either how much to import in the event of shortfall or optionally to export in case of surplus (Noureldinet *al.*, 2013). Rice is one of the most important crops in terms of human consumption (as opposed to animal feed) and is

Received 28 February, 2020

Accepted 30 June, 2020

Address Correspondence to:

yahayamohammed1986@gmail.com,

produced in 95 countries across the world (Maclean *et al.*, 2002). It is the staple food in many countries, accounting for more than 40% of global food production. People in the majority of countries in Asia depend on rice as their main source of nutrition, as well as for income and employment (Maclean *et al.*, 2002; Makino, 2011). Rice has been a major contribution to the financial income of many Asian countries as shown in the statistics, 50 % per annum is contributed to countries economy (Fahmiet *et al.*, 2013). Henceforth, numerous experiments were carried out to create a good quality paddy and to expand its yield due to increase demand in food. Due to consistent climate change some part of the paddy soil are prone to metal contamination (Sow *et al.*, 2013).

Reported in (Dahikaret *et al.*, 2014) is a study of basic requirements for applications of ANNs in yield prediction. Simple network architectures, with one hidden layer and back propagation of errors were tested for different predictors and crops, like cotton, sugarcane, wheat, rice and others. Soil parameters detected to be relevant for crop yield prediction were PH and concentrations of nitrogen, phosphate, potassium, organic carbon, calcium, magnesium, sulphur, manganese, copper and iron. In terms of atmospheric predictors, temperature, rainfall and humidity were the relevant features detected.

Monisha *et al.* (2005) developed a model for corn and soybean yield forecasting with climatic aspect by applying artificial neural network. They have considered the rainfall, Maryland corn and soybean yield data and predict the corn and soybean yield at state, regional and local levels by applying both the artificial neural network technique and the multiple linear regression model. Lastly they compared both the techniques and conclude that the ANN model gives more accurate yield prediction than the multiple linear regressions. (Saran *et al.*, 2014) proposed a new approach to estimate rice cultivation and harvest dates by using 8-day composite normalized difference vegetation index (NDVI) derived from Moderate Resolution Imaging Spectroradiometer (MODIS) data. In their work, they divided the rice growth state into 4 states, namely, nothing, growing, mature and harvest state and applied Hidden Markov Model (HMM). Then, they assigned the state to the NDVI time-series data

by using the Viterbi algorithm. By using those derived state, they were able to estimate the rice cultivation and harvest dates. The date estimation results were compared with the ground truth data to access the accuracy and they found that the average cultivation dates and harvest dates to have errors of 15.48 days and 6.525 days, respectively. They derived a simple linear regression model for wheat yield estimate and forecast based on NDVI images during the wheat grain filling period. They validated their results against official data and found good correlations between the two.

METHODOLOGY

Study Area and Data source

The data used in this research work, were collected from the Niger State bureau of statistics for the period of 16 years (2001 – 2016). Niger state with a population of 5,556,247 million people (National population commission, 2016) is located in the North central zone along the Middle Belt region of Nigeria. It is classified as one of the largest states in the country, spanning over 86,000 km² in land area with 80% of the land mass conducive for agriculture (Tolobonse, 2008).

Hidden Markov Model: A Hidden Markov Model (HMM) is a double stochastic process in which one of the stochastic processes is an underlying Markov chain which is called the hidden part of the model, the other stochastic processes is an observable one. Also a HMM can be considered as a stochastic process whose evolution is governed by an underlying discrete (Markov chain) with a finite number of state which are hidden, i.e. not directly observable (Enza, Daniele, 2007 and Lawal, 2017). Hidden Markov Model is characterized by the following

N = number of state in the model
 M = number of distinct observation symbols per state

Q = a state sequence of length T taking values from S ,

$$Q = q_1, q_2, q_3, \dots, q_T \quad (1)$$

O = an observation sequence consisting of T observations.

$$O = o_1, o_2, o_3, \dots, o_T \quad (2)$$

$A = \{a_{ij}\}$, a transition probability

matrix A , where each a_{ij} represents the

probability of moving from state S_i to state S_j ,

$$\sum_{j=1}^N a_{ij} = 1$$

with

$$B = \{b_j(o_t)\}, \text{ observation probability matrix}$$

Where

$b_j(o_t) = p(o_t | q_t = S_j)$ Is the probability that the symbol O_t is emitted when the system is in state S_j

If the observation is continuous a probability density function is used as follows:

$$\int_{-\infty}^{+\infty} b_j(x) dx = 1$$

$$\pi = \{\pi_j\}$$

An initial probability distribution, where π_i

Indicates the probability of starting in state S_i

.Also,

$$\sum_{i=1}^N \pi_i = 1.$$

The parameters of hidden of Markov model (HMM) denoted by

$$\lambda = (A, B, \pi) \tag{4}$$

Model Formulation:Hidden Markov Model is used to examine the impact of rainfall and temperature in forecasting rice yield. The quantity of rice yield to a great extent relies upon these climatic components they contribute colossally to rice yield in a growing years. But we can't ordinarily quantify how every one of them contribute to the general yield (the quantity of rice yield). The quantity of rice yield relies upon them and their measure are not static or deterministic but they vary randomly from year to year, which makes the measure of rice yield in each growing year to change.

This circumstance follows a doubly stochastic process, with the measure of rice yield per hectare as the observation of the HMM depending on the state (measure of rainfall and temperature). In perspective on this, we have taken the measure of rice yield per hectare within a growing year as an emission of the HMM while the measure of rainfall and temperature within a similar period is taken as state of our model. We make the following assumptions.

1. The transition between the states is governed by first order Markov reliance as represented by equation (5)

$$P\{X_{n+1} = j | X_0 = i_0, \dots, X_{n-2} = i_{n-2}, X_{n-1} = i_{n-1}, X_n = i\} = P_{ij} \tag{5}$$

2. The probability of conveyance of generating current observation symbol depends on current state, as represented by equation (6)

$$P(O | Q, \lambda) = \prod_{t=1}^T P(o_t | q_t, \lambda) \tag{6}$$

3. Quantity of rainfall is viewed as low in the event that it is under 193.65mm
4. Quantity of rainfall is viewed as high in the event that it is above 194.65mm
5. Measure of temperature is viewed as low in the event that it is underneath 27.95⁰c
6. Measure of temperature is viewed as high in the event that it is above 28.95⁰c
7. Quantity of rice yield is viewed as low on the off chance that it is underneath 2.986 metric tons
8. Quantity of rice yield is viewed as high in the event that it is above 3.992 metric tons
9. Quantity of rice yield is viewed as moderate in the event that it is in range of (2.986-3.992) metric tons

Following (Lawal, 2017), we have the following states and observations for our model

State 1: low rainfall and low temperature

State 2: low rainfall and high temperature

State 3: high rainfall and low temperature

State 4: high rainfall and high temperature

Observations.

The classification of states and the observations, and the assumption made in this work are based on the study area and the data obtained




Figure 1: Transition Diagram of the Rice Yield Model

Making Prediction with the Model: Making prediction with the model is done with the training of parameters of the model. The parameters of the model are initialized then trained using Baum-Welch algorithm to attend maximum likelihood. The forward probability of the training observation sequence is determined from time $t=1$ to time T using Forward Algorithm (Rabiner, 1989 and Lawal, 2017). To predict the next state at time $T+1$ and its observation given the present state at time T , we calculate forward probability for every conceivable observation of the states, at that point, the sequence with most elevated value of the forward probability at time $T+1$ is taken as predicted state and its observation. The prediction is made for the following three years (year 2 at time $T+2$, year 3 at time $T+3$ and year 4 at time $T+4$).

RESULTS AND DISCUSSION

Application of the Hidden Markov Model for Rice Yields Forecast: The data used in this illustration was collected from the archive of Niger State Bureau of Statistics for the period of 16 years (2001-2016). The summary of the data is presented in Table 1

Table 1: The States and Observations of the Hidden Markov Model for Period of sixteen years

Years	States	Observations
2001	1	L
2002	1	M
2003	2	M
2004	2	M
2005	2	M
2006	4	M
2007	3	H
2008	3	H
2009	1	H
2010	4	H
2011	2	H
2012	2	M
2013	2	H
2014	4	H
2015	3	H
2016	4	H

Validity Test for the Model: To test for the validity of the model, we divide the data set into two sets, one of the sets was used to build the test model (HMM1) and the other set was used to test for the reliability of the model. We estimate the parameters of HMM1 utilizing Rainfall, Temperature and Rice yield data from 2001 to 2012, then test for the reliability of the model using data set for the following years 2013, 2014, 2015 and 2016.

The Transition Count Matrix, Pseudo count Transition Matrix and Transition Probability Matrix are given in Equations (7), (8) and (9) respectively.

$$C = \begin{bmatrix} 1 & 1 & 0 & 1 \\ 0 & 3 & 0 & 1 \\ 1 & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 \end{bmatrix} \quad (7)$$

$$S = \begin{bmatrix} 2 & 2 & 1 & 2 \\ 1 & 4 & 1 & 2 \\ 2 & 1 & 2 & 1 \\ 1 & 2 & 2 & 1 \end{bmatrix} \quad (8)$$

$$A = \begin{bmatrix} 0.2857 & 0.2857 & 0.1428 & 0.2857 \\ 0.1250 & 0.5000 & 0.1250 & 0.2500 \\ 0.3333 & 0.1666 & 0.3333 & 0.1666 \\ 0.1666 & 0.3333 & 0.3333 & 0.1666 \end{bmatrix} \quad (9)$$

While Observation count matrix, Pseudo count Observation matrix and Observation probability matrix are given in equations (10), (11) and (12), respectively.

$$E = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 4 & 1 \\ 0 & 0 & 2 \\ 0 & 1 & 1 \end{bmatrix} \quad (10)$$

$$D = \begin{bmatrix} 2 & 2 & 2 \\ 1 & 5 & 2 \\ 1 & 1 & 3 \\ 1 & 2 & 2 \end{bmatrix} \quad (11)$$

$$B = \begin{bmatrix} 0.4000 & 0.2000 & 0.2222 \\ 0.2000 & 0.5000 & 0.2222 \\ 0.2000 & 0.1000 & 0.3333 \\ 0.2000 & 0.2000 & 0.2222 \end{bmatrix} \quad (12)$$

The initial state probability is given below

$$\pi = [0.25, 0.4167, 0.1666, 0.1666] \quad (13)$$

The general HMM1 is represented by equation (14)

$$\lambda_1 = (A, B, \pi) \quad (14)$$

After 1000 iteration of Baum Welch Algorithm, equation (14) settled to (15)

$$\lambda_1^* = (\hat{A}, \hat{B}, \hat{\pi}) \quad (15)$$


When

$$\hat{A} = \begin{bmatrix} 0.0000 & 1.0000 & 0.0000 & 0.0000 \\ 0.0000 & 0.4283 & 0.0000 & 0.5717 \\ 0.0000 & 0.0000 & 1.0000 & 0.0000 \\ 0.0000 & 0.0000 & 0.5717 & 0.4283 \end{bmatrix} \quad (16)$$


$$\hat{B} = \begin{bmatrix} 1.0000 & 0.0000 & 0.0000 \\ 0.0000 & 1.0000 & 0.0000 \\ 0.0000 & 0.0000 & 1.0000 \\ 0.0000 & 0.2102 & 0.7898 \end{bmatrix} \quad (17)$$

Comparison of the Predicted States and Observations, and the Actual States and Observations from the Dataset

Predicted States and Observations:



Actual States and Observations:



The parameter of the HMM1 were determined utilizing rainfall, temperature and rice yield data from 2001 to 2012. After 1000 iterations of the Baum Welch Algorithm, λ_1 settled to another model, λ_1^* , this new model was used to test for rice yield for 2013, 2014, 2015, and 2016. From the test, the HMM1 was in state 2 at time T (2012) emitting Moderate rice yield, at that point, it then make move to state 2 at time T+1 (2013) emitting a Moderate rice yield, then make move to state 4 at time T+2(2014) emitting a High rice yield, next, it make move to state 3 at time T+3(2015) emitting a High rice yield and move to state 3 at time T+4 (2016) emitting a High rice yield. The transition between the states are governed by the first order Markov dependence as referenced in the past section. The validity test shows 80% precision in the rice forecast when compared with the actual rice yield from the dataset.

Hidden Markov Model (HMM2) for future forecast: HMM2 was developed to forecast rice yield for future years, the parameters of the HMM2 were determined utilizing rainfall, temperature and rice yield data from 2001 to 2016, at the point, we made forecast for 2017, 2018, 2019, and 2020.

Transition Count Matrix

$$C = \begin{bmatrix} 1 & 1 & 0 & 0 \\ 0 & 4 & 0 & 2 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 2 & 0 \end{bmatrix} \quad (18)$$

Pseudo count Transition Matrix

$$S = \begin{bmatrix} 2 & 2 & 1 & 1 \\ 1 & 5 & 1 & 3 \\ 2 & 1 & 2 & 2 \\ 1 & 2 & 3 & 1 \end{bmatrix} \quad (19)$$

Transition Probability Matrix

$$A = \begin{bmatrix} 0.2857 & 0.2857 & 0.1428 & 0.2857 \\ 0.1000 & 0.5000 & 0.1000 & 0.3000 \\ 0.2857 & 0.1428 & 0.2857 & 0.2857 \\ 0.1428 & 0.2857 & 0.4285 & 0.1428 \end{bmatrix} \quad (20)$$

Observation Count Matrix

$$C = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 4 & 2 \\ 0 & 0 & 3 \\ 0 & 1 & 3 \end{bmatrix} \quad (21)$$

Pseudo count Observation Matrix

$$S = \begin{bmatrix} 2 & 2 & 2 \\ 1 & 5 & 3 \\ 1 & 1 & 4 \\ 1 & 1 & 4 \end{bmatrix} \quad (22)$$

Observation Probability Matrix

$$B = \begin{bmatrix} 0.4000 & 0.2222 & 0.1535 \\ 0.2000 & 0.5555 & 0.2307 \\ 0.2000 & 0.1111 & 0.3076 \\ 0.2000 & 0.1111 & 0.3076 \end{bmatrix} \quad (23)$$

Initial State Probability

$$\pi = [0.1875 \quad 0.1875 \quad 0.250 \quad 0.375] \quad (24)$$

$$\lambda_2 = (A, B, \pi) \quad (25)$$

After 1000 iteration of Baum Welch Algorithm, equation (26) stabilized to (27)

$$\lambda^* = (\hat{A}, \hat{B}, \hat{\pi}) \quad (28)$$

Where

$$\hat{A} = \begin{bmatrix} 0.0000 & 1.0000 & 0.0000 & 0.0000 \\ 0.0000 & 0.7500 & 0.0000 & 0.2500 \\ 0.0000 & 0.0000 & 0.8750 & 0.1250 \\ 0.0000 & 0.0000 & 1.0000 & 0.0000 \end{bmatrix} \quad (27)$$

$$\hat{B} = \begin{bmatrix} 1.0000 & 0.0000 & 0.0000 \\ 0.0000 & 1.0000 & 0.0000 \\ 0.0000 & 0.0000 & 1.0000 \\ 0.0000 & 1.0000 & 0.0000 \end{bmatrix} \quad (28)$$

Future Forecast States and Observations:

States: 4(2016) → 3(2017) → 3(2018) → 3(2019) → 3(2020)
 Observations: H H H H H

The parameter of the HMM2 were determined using rainfall, temperature and rice yield data from 2001 to 2016. After 1000 iteration of the Baum Welch algorithm, λ_2 , settled to another model, λ_2^* , this new model was then used to make a forecast for future years. From the forecast, the HMM2 was in state 4 at time T (2016) emitting High rice yield, at that point, it then make move to state 3 at time T+1 (2017) emitting High rice yield. Similar interpretation is given to move to state 3 at time T+2 (2018), move to state 3 at time T+3 (2019), move to state 3 at time T+4 (2020) all emitting High rice yield.

Conclusion

A hidden Markov Model to forecast Rice yield with respect to major climatic elements (rainfall and temperature) has been developed and implemented in Niger State. The validity test for the model show 80% precision in Rice yield forecast. Both the validity test and the future forecast shows pervasiveness of high Rice yield, this shows that Niger State is a better place for rice farming. The model could likewise be used to forecast Yield of other crops with little or no modification.

REFERENCES

- Ajith, K., V. Geethalakshmi, K.P. Ragunath, S. Pazhanivelan and Dheebakaran Ga. 2017. Rice Yield Prediction Using MODIS – NDVI (MOD13Q1) and Land Based Observations. *Int.JCurr.Microbiol. App.Sci.*6(12):22772293.doi:https://doi.org/10.20546/ijcmas.2017.612.263.
- Ayanwale, F.S., Akinyosoye, V. S. A. And Oni, A. O. 2011. Supply Response in Nigeria: Winter Change. Central Bank of Nigeria Monthly Bulletin, 2011.
- Akande, T. (2003). The rice sector in Nigeria. United Nation Crop Project (UNCP). Country agriculture project on trade liberalization in agricultural sector and the environment, Geneva.10pp.
- Ahn, P.M. (1993). Tropical soils and Fertiliser Use, London: Longman Scientific & Technical.
- Abubakaret al (2014). Markov Chain Model and its Application to Annual Rainfall Distribution for Crop Production. *American Journal of Theoretical and Applied Statistics*.Vol.3, No2, 2014, pp. 39-43.doi:10.11648/j.ajtas.20140302.12.
- Anderson, T.W., and L.A. Goodman. “Statistical Inference about Markov Chains.” *Annals of Mathematical Statistics*, 28(1) (1957): 89-110.
- Agrawal, R. and Jain, R.C. (1996). Forecast of sugarcane yield using eye estimate along with plant characters, *Biom. J.*, 38, 731 39.
- Akintayo, O.I., &Radji, M. (2012). Determinants of yield gap in upland rice production in South-western Nigeria. *International Journal of Agriculture and Rural Development*, 23(3),1595- 9716.
- Abramowitz, M. and Stegun, I.A. (1965). Hand book of mathematical functions. Washington, D.C.: National Bureau of Standards.
- Akande, S.O. (2003). The Nigerian Rice Economy in a Complete World: Constraints, Opportunities and Strategic Choices: Rice Production Systems in Nigeria: A Survey. Abidjan: Côte d’Ivoire, West Africa Rice Development Association (WARDA).
- Basso, B., Cammarano, D. and Carfagna, E. 2013.Review of crop yield forecasting methods and early warning systems. Proceedings of the First Meeting of the Scientific Advisory Committee of the Global Strategy to Improve Agricultural and Rural Statistics, FAO Headquarters, Rome, Italy, 18–19 July.
- Bastiaanssen WGM, Ali S, 2003. A new crop yield forecasting model based on satellite measurements applied across the Indus Basin, Pakistan. *Agr Eco Environ* 94(3): 321340.http://dx.doi.org/10.1016/S0167-8809 (02)00034-8
- Doraiswamy, P.C., Moulin, S., Cook, P.W., Stern, A., 2003. Crop Yield Assessment from Remote Sensing. *Photogrammetric Engineering & Remote Sensing* 69, 665–674.

- Dahikar MSS, Rode SV. Agricultural crop yield prediction using artificial neural network approach. *International Journal of Innovative Research in Electrical, Electronics, Instrumentation and Control Engineering (IJIREICE)*. 2014 Jan; 2(1):1–4.
- Daramola, B. (2005). Government policies and competitiveness of Nigeria rice economy. Proceeding of a workshop on rice policy and food security in Sub-Saharan Africa organized by WARDA, Cotonou. Republic of Benin Nov. 07-09, 2005.
- Ejiogu, A.O & Okoli, V.B.N (2012). Bird scaring technologies in rice production: The need for policies prohibiting participation of women and children. *Global Advanced Research Journal of Food Science and Technology*, 1(3), 031-038.
- Emodi, I. A. and Madukwe, M. C. (2008). A Review of policy, Acts and Initiatives in Rice Innovation System in Nigeria. *Journal of Agricultural Extension*. 12 (2): 12 – 21.
- Food and Agriculture Organization (FAO) (2010). FAO Agricultural Bulletin of Statistics 3(2): 15. Rome, Italy. www.fao.org.
- Enza, Messina; Daniele, Toscani (2007). Hidden Markov models for scenario generation, *IMA Journal of Management Mathematics* 19: 379-401.
- Feng G, Cobb S, Abdo Z, Fisher DL, Ouyang Y, Adeli A and Jenkins JN (2016) Trend analysis and forecast of precipitation, reference evapotranspiration, and rainfall deficit in the Blackland Prairie of eastern Mississippi. *Journal of Applied Meteorology and Climatology* 55, 1425–1439.
- Fahmi, Z., Abu Samah, B., & Abdullah, H. (2013). Paddy industry and paddy farmers' well-being: A success recipe for agriculture industry in Malaysia. *Asian Social Science*, 9(3), 177–181. <https://doi.org/10.5539/ass.v9n3p177>.
- FMARD – Federal Ministry of Agriculture and Rural Development 2001.
- Olorunfemi, O.O. (2006). Technical efficiency differential in rice production technologies in Nigeria. AERC Research paper 154 African Economic Research Consortium, Nairobi.
- Gagniuc, Paul A. (2017). Markov Chains: From Theory to Implementation and Experimentation. USA, NJ: John Wiley & Sons. pp. 1–235. [ISBN 978-1-119-38755-8](https://doi.org/10.1002/9781119387558).
- Gobin A, Kersebaum KC, Eitzinger J, Trnka M, Hlavinka P, Takáč J, Kroes J, Ventrella D, Dalla Marta A, Deelstra J, Lalić B, Nejedlik P, Orlandini S, Peltonen-Sainio P, Rajala A, Saue T, Şaylan L, Stričević R, Vučetić V and Zoumides C (201) Variability in the water footprint of arable crop production across European regions. *Water* 9, 93, 10.3390/w902009.
- Huang J, Wang X, Li X, Tian H, Pan Z, 2013. Remotely sensed rice yield prediction using multi-temporal NDVI data derived from NOAA's AVHRR. *PLOS ONE* 8(8):e70816. <http://dx.doi.org/10.1371/journal.pone.0070816>
- Imolehin, E. D. and Wada, A. C. (2000). Meeting the Rice Production and Consumption Demands of Nigeria with Improved Technologies. *International Rice Commission Newsletter*. 49: 33–41.
- Isaac, B.D., Irene, S.E. 2014. Factors that influence household demand for locally produced Brown Rice in Ghana. *Journal of Economics and Sustainable Development*. 5(7) 14 - 24
- IPCC, 2014. Climate Change: Synthesis Report. IPCC Fifth Assessment Synthesis Report; Intergovernmental Panel on Climate Change, Geneva, Switzerland, 151pp. <http://www.ipcc.ch/report/ar5/syr/>. [2 Nov 2014].
- Jain, R.C. and Agrawal, R. (1992). Probability model for crop yield forecasting, *Biom. J.*, 34, 501–511.
- Kassali, R., Kareem, R.O., Oluwasola, O., and Ohaegbunan, O.M. 2010. Analysis of Demand for Rice in Ile-Ife, Osun State, *Nigeria Journal Sustainable Development in Africa*, 12(2).

- Kogan, F., Salazar, L., Roytman, L., 2012. Forecasting crop production using satellite-based vegetation health indices in Kansas, USA. *International Journal of Remote Sensing* 33,27982814
- Kogan, F., Kussul, N., Adamenko, T., Skakun, S., Kravchenko, O., Kryvobok, O., Shelestov, O., Kolotii, A., Kussul, O., Lavrenyuk, A., 2013. Winter wheat yield forecasting in Ukraine based on Earth observation, meteorological data and biophysical models. *International Journal of Applied Earth Observation and Geoinformation* 23, 192–203.
- Kouadio, L., Newlands, N. K., Davidson, A., Zhang Y. and Chipanshi, A. 2014. Assessing the Performance of MODIS NDVI and EVI for seasonal crop yield forecasting at the eco district scale. *Agric. For. Meteorol.* 173: 74– 84
- Lopresti, M. F., Di Bella, C. M., Degioanni, A. J. 2015. Relationship between MODIS-NDVI data and wheat yield: A case study in Northern Buenos Aires province, Argentina, *Information processing in Agriculture*. 2: 73– 84.
- Lawal A. (2017). Stochastic model of rainfall precipitations for crop production in some selected states of north central Nigeria. Unpublished PhD thesis, Department of Mathematics Federal University of Technology Minna.
- Makino, A. (2011). Photosynthesis, grain yield, and nitrogen utilization in rice and wheat. *Plant Physiology*, 155(1), 125– 129.
- Mahboob, M. G., Islam, A. T. and Deshapriya, L. 2016. Rice mapping and monitoring in Sylhet region of Bangladesh using MODIS NDVI. *Proceedings of Asia Flux Mini-Workshop on remote sensing and ecological /environmental monitoring*, March 2016, Thailand, 26- 30.
- Manjunatha, M., Parkavi, A. (2018). Estimation of Arecanut Yield in Various Climatic Zones of Karnataka using Data Mining Technique: A Survey. 2018 *International Conference on Current Trends towards Con- verging Technologies (ICCTCT)*.
- Marlia, M., Nasuddin, O. &Fazleen, A. 2011. Determination of Consumers Purchasing Behaviour for Rice in Malaysia. *American International Journal of Contemporary Research*.1 (3) 159 – 167.
- Matis, J.H., Saito, T., Grant, W.F. Iwig, W.C. and Ritchie, J.T. (1985). A Markov chain approach to crop yield forecasting, *Agricultural Systems*, 18, 171 87.
- Monisha Kaul M, Robert L, Hill H, Walthall C. Artificial neural networks for corn and Soybean yield prediction, Elsevier. *Agricultural System*. 2005; 85(1):1–18.
- Matis, J.H., Birkett, T. and Boudreaux, D. (1989). An application of the Markov chain Approach to forecasting cotton yield from surveys, *Agricultural Systems*, 29, 357 70.
- Mostafa, K. K. M. 2015. Use of GIS and remote sensing in mapping rice areas and forecasting its production at large geographical extent. Ph. D. thesis. University of Calgary, Alberta, Canada.
- Miguel-Ayala L, van Eupen M, Zhang GP, Pérez-Soba M, Martorano LG, Lisboa LS and Beltrao NE (2016) Impact of agricultural expansion on water footprint in the Amazon under climate change scenarios. *Science of the Total Environment* 569–570, 1159–1173.
- Maclean, J. L., Dawe, D. C., Hardy, B. & Hettel, G. P. (2002). *Rice almanac source book for the most important economic activity on earth*. Published by CABI (Oxon, UK) for IRRI (Manila) WARDA (Cote d'Ivoire), CIAT (Colombia) and FAO (Rome, Italy).
- Nuarsa IW, Nishio, F, Hongo C, 2012. Rice yield estimation using Landsat ETM+ data and field observation. *J AgricSci* 4(3): 45- 56.
- Noureldin NA, Aboelghar MA, Saady HS, Ali AM, 2013. Rice yield forecasting models using satellite imagery in Egypt. *Egypt J Remote SensSpace Sci* 16(1): 125-131. <http://dx.doi.org/10.1016/j.ejrs.2013.04.005>.
- Osman J, Inglada J, Dejoux JF. Assessment of a Markov logic model of crop rotations for early crop mapping, Elsevier. *Computers and Electronics in Agriculture*. 2015; 113:234–43.

- Onu D O, Obike K. C., Ebe F. C. and Okpara B. O. 2015. Empirical Assessment of the Trend in Rice Production and Import in Nigeria (1980 – 2013): *International Journal of Agricultural Science*. 5 (6) 150 – 158
- Oliver K (2009). Department of mathematics, Harvard University Cambridge, MA 02138, USA
- Oko, A.O and Ugwu, S.I. (2011). The proximate and mineral composition of five major Rice varieties in Abakaliki, South-Eastern Nigeria. *International journal of plant physiology and Biochemistry* 3(2): 25-27.
- Parzen. Emmanuel (1962). Stochastic processes, Oakland CA: Holden Day
- Ramasubramanian, V. and Jain, R.C. (1999). Use of growth indices in Markov chain models for crop yield forecasting, *Biom. J.*, 41, 99 109.
- Ramasubramanian V., Agrawal, R. and Bhar, L. (2010). Crop forecasting using multiple Markov chains, *Assam Stat. Rev.*, 24(1), 37-56.
- Ross S.M. (1989). Introduction to probability Model Academic Press, Ltd, London.
- Rabiner, L.R. (1989). A tutorial on hidden Markov models and selected applications in speech recognition. *Proc. IEEE*, 77, .257-286.
- Son NT, Chen CF, Chen CR, Chang LY, Duc HN, Nguyen LD, 2013. Prediction of rice crop yield using MODIS EVI-LAI data in the Mekong Delta, Vietnam. *Int J Remote Sens* 34(20): 7275-7292. <http://dx.doi.org/10.1080/01431161.2013.818258>.
- Singh, R. and Ibrahim, A.E.I. (1996). Use of spectral data in Markov chain model for crop yield forecasting. *Jour. Ind. Soc. Remote Sensing*, 24, 145 52.
- Sow, A. Y., Ismail, A., &Zulkifli, S. Z. (2013). Geofractionation of heavy metals and application of indices for pollution prediction in paddy field soil of Tumpat, Malaysia. *Environmental Science and Pollution Research*, 20(12), 8964–8973. <https://doi.org/10.1007/s11356-013-1857-9>.
- Saran. S, Teerasit. K., Kitti .C. Preesan .R. & Thitiporn C, Itsuo K. (2014). Rice Cultivation and Harvest Date Estimation Using MODIS NDVI Time-series. Data Nagatsuta-cho, Midori-ku, Yokohama Japan, 226-8503.
- Tologbonse, D. (2008 December). Policy Issues In Meeting Rice Farmers Agricultural Information Needs in Niger State. *Journal of Agricultural Extension*. 12:2:84-94.
- Tonifelix, C .M. 2017. The New Writer of Abakaliki Local Rice, 8(6): 2 -8.
- Veenadhari S, Mishra B, Singh CD. Soybean Productivity Modelling using Decision Tree Algorithms. *International Journal of Computer Applications*. 2011; 27(7):975–8887.
- WARDA (1996). ANNUAL Report. West African Rice Development Association M'be, Cote D'Ivoire.
- WARDA (2005). Rice policy and food security in Sub-sahara Africa. Proceeding of a workshop held on 7 – 9 Nov.2005, Cotonou, Benin.
- Y. Shen, L. Di, L. Wu, G. Yu, H. Tang & G. Yu, (2012) "Hidden Markov Models for corn progress percents estimation in multivariate time series," in *Agro-Geoinformatics (Agro-Geoinformatics)*, 2012 *First International Conference on*, pp. 1-6.
- Zhuo L, Mekonnen MM and Hoekstra AY (2016) the effect of inter-annual variability of consumption, production, trade and climate on crop-related green and blue water footprints and inter-regional virtual water trade: a study for China (1978–2008). *Water Research* 94, 73 –85.
- Zoumidis C, Bruggeman A, Hadjikakou M and Zachariadis T (2014) Policy-relevant indicators for semi-arid nations: the water footprint of crop production and supply utilization of Cyprus. *Ecologic-al Indicators* 43, 205–2.